



# Similarities and Differences Between Bipolar and Depression

by BECKY WICKS

---

## Bipolar vs Depression

Being diagnosed with a mental illness can feel like the end of the world. It may make you feel like you are completely alone, or it may in fact make you feel as though you finally have some answers to the way you have been feeling.

It is important to realize that mental health diagnoses are not always cut and dried, and sometimes there may be crossovers in your symptoms, thus meaning a condition is harder to diagnose. Both depression and bipolar sit on the mental health spectrum, and have many similar symptoms, meaning at times they can be hard to distinguish from one another.

Depression is one of the most commonly diagnosed mental health issues, and is recognizable by symptoms like feelings of guilt, worthlessness or hopelessness, issues with sleeping patterns (sleeping too much or too little), a general sense of apathy and lack of drive to do anything, and an overall low mood.

Bipolar spectrum disorders involve depressive episodes, but also entail manic episodes. Manic episodes are the main distinguishing factor between unipolar depression and bipolar disorder, and are categorized by symptoms such as an elevated mood and ego, grandiose thinking and behaviour, impulsive behaviour such as overspending and risky sexual behaviour, and rapid thoughts.

Bipolar disorder is actually a spectrum with more than one disorder sitting on it; bipolar II is a less severe form of bipolar, and is recognizable by longer periods of depression, and shorter periods of hypomania — often mistaken for just a really good mood.

In contrast to this, bipolar I is a more intense version of the disorder, categorized by longer periods of full mania. This is a more intrusive form of mania, often progressing on from hypomania, and often results in the need for inpatient treatment.

One type of psychiatric symptoms that can affect both certain types of bipolar and depression is that of psychosis — it is possible to be diagnosed with psychotic depression, and in the more intense bipolar I patients may experience psychotic symptoms at the very heights of mania or the lowest points of depression.

Psychosis can be a very distressing experience, and may include symptoms such as auditory or visual hallucinations, delusions such as the belief that someone has bad intentions that you have special powers, or severe paranoia.

## Misdiagnosis

Misdiagnosis of bipolar as depression is very common, as people with bipolar often exhibit only symptoms of depression at first, or the depressive symptoms are the only ones they feel they need to seek help for.

---

This is especially prevalent in patients with bipolar II patients, as they may feel that the hypomania is a relief from the low moods, or that they are simply having a better time than they usually are. The risk here is eventually falling into a more intense and damaging manic episode as you may not have the coping skills in place, or may not be able to recognize the more dangerous symptoms setting in.

More and more people with depression are now being re-diagnosed as having bipolar disorder as their illness progresses, or as doctors have a better insight into the patient's history.

It is also possible that the traditional treatment of antidepressants given to patients diagnosed with depression may in fact trigger underlying bipolar disorder. Antidepressant-induced mania is very much a legitimate risk, and therefore when you visit a doctor in order to ask for help, you must be completely honest about all your symptoms — even ones that may seem like positive things at the time.

A diagnosis of a mental health issue is based on multiple factors, but two of the most important things that a professional will rely on is observation of current behaviours, and history, impacting the need for honesty even further.

*Next page: how each condition is treated.*

## **Treatments for Each Condition**

Whilst there is some crossover in treatment for the two disorders (e.g. antidepressants may be used in conjunction with other medications for bipolar treatment), there are some important additions to the treatment of bipolar disorder.

Whilst depression tends to be (traditionally) treated with antidepressants such as Prozac and Zoloft, bipolar disorder may also be treated with mood stabilizers such as Lamictal and Epilim, and if psychotic symptoms have been identified then there may also be the addition of antipsychotics such as Abilify or Quetiapine.

Medication is an important part of treatment in stabilizing both disorders, but it is vital that chemical treatment is strengthened by the addition of talking therapies. There are many options available for talking therapy, with options ranging from counselling to psychotherapy.

In more extreme cases of both depression and bipolar disorder, there may be a need for a treatment known as ECT (electroconvulsive therapy), a procedure by which seizures are induced in the brain in order to provide relief from more treatment resistant symptoms. This is usually only used in very intense cases however, where many medications have not worked.

There are many lifestyle changes that can be made to help the symptoms of mental illness, and many of them apply to both depression and bipolar. For example, it is important to maintain a good sleep routine as best you can. Going to bed at the same time and getting up at the same time will have a positive impact on your mental health, as will trying to stick to a basic routine throughout your daily tasks.

It is important to avoid street drugs and alcohol as these will not help the situation in the long run, and will eventually exacerbate any symptoms, or may cause a full relapse. Gentle exercise such as walking or yoga can help to raise serotonin levels, and performing activities such as meditation and mindfulness can help to reconnect the mind with the body, thus causing greater self-awareness.

Whilst it is important to monitor moods, especially through a self kept mood diary, it is also important to not be overly hyper vigilant of moods, as this may lead to feelings of negativity surrounding any kind of good time, as they may then be connected with anxiety and worry.

If you feel that you may be experiencing some mental health related symptoms, you should try and seek help at the earliest opportunity. It has been proven across multiple studies that early intervention is helpful to

---

achieving the best possible outcome with respect to a mental health diagnosis.

Seeking help early will mean you can build coping strategies, and will be able to recognize warning signs in yourself before a crisis hits. Reaching out to friends and family is also something to do in times of mental health distress as they can provide valuable help and support during difficult times.

In times of crisis, it is important to reach out to a mental health professional as soon as possible so that the correct help can be accessed. If the crisis is severe and happens outside of the operating hours of a standard mental health team then a visit to hospital is an absolute necessity.